

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460



OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

30 September 2014

VIA EMAIL

Ms. Michelle Sharpe-Kass
Product Registration Manager
BASF Corporation
26 Davis Drive
P.O. Box 13528
Research Triangle Park, NC 27709

Subject: Label Amendment: Incorporating Supplemental Labels into Master Label
Product Name: Priaxor® Xemium® Brand Fungicide
EPA Registration Number: 7969-311
Application Date: April 8, 2014
Decision Number: 490477

Dear Ms. Sharpe-Kass:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

SEE NEXT PAGE

with FIFRA section 6(e). If you have any questions, please contact Michael Walsh by phone at (703) 308-2972 or via email at "walsh.michael@epa.gov".

Sincerely,


A handwritten signature in black ink that reads "Mark ES". The "M" is large and stylized, followed by "ark" in a smaller, cursive script. To the right, the letters "ES" are written in a large, bold, cursive font.

Mark Suarez, Product Manager 13
Insecticide Branch
Office of Pesticide Programs

Enclosure

Priaxor®

Xemium® Brand Fungicide



For disease control and plant health in the following crops: barley, berries and small fruits, Brassica leafy vegetables, bulb vegetables, corn (all types), cucurbit vegetables, dried shelled peas and beans, edible-podded legume vegetables, fruiting vegetables (including tomato), grapes, leafy vegetables, oats, oilseed crops (including flax seed, rapeseed, safflower, and sunflower), peanut, root vegetables, rye, sorghum and millet, soybean, strawberries, succulent shelled peas and beans, sugar beet, sugarcane, tree nuts, tuberous and corm vegetables (including potato), wheat and triticale

Powered by **Xemium®** and **F500®** fungicides

Active Ingredients:

fluxapyroxad*: 1 <i>H</i> -Pyrazole-4-carboxamide, 3-(difluoromethyl)-1-methyl-N-(3',4',5'-trifluoro[1,1'-biphenyl]-2-yl)-	14.33%
pyraclostrobin**: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1 <i>H</i> -pyrazol-3-yl]oxy]methyl]phenyl]methoxy-, methyl ester).	28.58%

Other Ingredients: 57.09%

Total: 100.00%

* Equivalent to 1.39 pounds of fluxapyroxad per gallon

** Equivalent to 2.78 pounds of pyraclostrobin per gallon

EPA Reg. No. 7969-311

EPA Est. No.

**KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Shake Well Before Using

Net Contents:

BASF Corporation
26 Davis Drive, Research Triangle Park, NC 27709

FIRST AID

If swallowed

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- **DO NOT** induce vomiting unless told to do so by a poison control center or doctor.
- **DO NOT** give anything to an unconscious person.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed. Avoid contact with skin or clothing.

Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. For more options, refer to **Category A** on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high

water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater.

Surface Water Advisory

This product is classified as having high potential for reaching aquatic sediment via runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this active ingredient or its degradates from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

Groundwater Advisory

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves (made of any waterproof material)
- Shoes plus socks

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. **DO NOT** store near food or feed.

Pesticide Disposal

Wastes resulting from using this product may be disposed of on-site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representatives at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity ≤ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

STORAGE AND DISPOSAL *(continued)*

Container Handling *(continued)*

Triple rinse containers too large to shake (capacity > 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable Container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

Triple rinse as follows: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

When this container is empty, replace the cap and seal all openings that have been opened during use; return the container to the point of purchase or to a designated location. This container must only be refilled with a pesticide product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transport. **DO NOT** transport if this container is damaged or leaking. If the container is damaged, or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling, if available, or dispose of container in compliance with state and local regulations.

(continued)

In Case of Emergency

In case of large-scale spillage regarding this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation 1-800-832-HELP (4357)

Steps to be taken in case material is released or spilled:

- In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.
- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

Product Information

This package contains **Priaxor® Xemium® brand fungicide**, a suspension concentrate (SC) containing the active ingredients fluxapyroxad and pyraclostrobin. The active ingredients in **Priaxor** belong to two classes of fungicides, the strobilurins or Quinone Outside Inhibitors (QoI) and the succinate-dehydrogenase (SDH) inhibitor classes. To maximize disease control, apply **Priaxor** in a regularly scheduled protective spray program and use in a rotation program with other fungicides.

Preventive applications optimize disease control, resulting in improved plant health. The increase in plant health comes from the combined effect of disease control (including fungal diseases listed in Crop-specific directions), improved growth efficiency and improved stress tolerance. Overall increased plant health may result in an improvement in crop growth and crop quality as well as increased crop yields.

Because of its high specific activity, **Priaxor** has good residual activity against target fungi.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>.

Priaxor is not for use in greenhouse or transplant production.

Modes of Action

Fluxapyroxad and pyraclostrobin, the active ingredients of **Priaxor**, belong to the groups of respiration inhibitors classified by the U.S. EPA and Canada PMRA as target site of action **Group 7** and **Group 11** fungicides, respectively.

Resistance Management

Priaxor contains fluxapyroxad and pyraclostrobin, a premix of a **Group 7** and a **Group 11** fungicide, and is effective against pathogens resistant to fungicides with modes of action different from those of target site **Group 7** and **Group 11**, such as dicarboximides, sterol inhibitors, benzimidazoles, or phenylamides. Fungal isolates resistant to **Group 7** or **Group 11** fungicides may eventually dominate the fungal population if **Group 7** or **Group 11** fungicides are used predominantly and repeatedly in the same field in successive years as the primary method of control for the targeted pathogen species, especially if resistance to either **Group 7** or **Group 11** fungicides is already present in the pathogen population. This may result in reduction of disease control by **Priaxor** or other **Group 7** or **Group 11** fungicides. To maintain the performance of **Priaxor** in the field, **DO NOT** exceed the specified number of sequential applications of **Priaxor** or the total number of applications of **Priaxor** per season stated in **Table 1. Priaxor® Xemium® brand fungicide Restrictions and Limitations Overview** and **Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions**. Adhere to the label instructions regarding the sequential use of **Priaxor** or other target site of action **Group 7** and **Group 11** fungicides that have a similar site of action on the same pathogens.

Resistance Management Advisory

The following recommendations may be considered to delay the development of fungicide resistance:

1. **Tank mixtures** - **Priaxor** provides more effective resistance management of most of its target pathogens, because it is a premix of two fungicides with different modes of action. If **Priaxor** is used in tank mixtures with fungicides from different target site of action groups that are registered/permitted for the same use and that are effective against the pathogens of concern, use at least the minimum labeled rates of each fungicide in the tank mix. For tank mix exceptions, see **Additives and Tank Mixing Information** section and **Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions**.
2. **IPM** - Integrate **Priaxor** into an overall disease and pest management program. Follow cultural practices known to reduce disease development. Consult your local extension specialist, crop advisor and/or BASF representative for additional IPM strategies established for your area. **Priaxor** may be used in agricultural extension advisory (disease forecasting) programs, which recommend application timing based on environmental factors favorable for disease development.
3. **Monitoring** - Monitor efficacy of all fungicides used in the disease management program against the targeted pathogen and record other factors that may influence fungicide performance and/or disease development. If a **Group 7** or **Group 11** target site fungicide such as **Priaxor** appears to be less or no longer effective against a pathogen that it previously controlled or suppressed, contact a BASF representative, local extension specialist, or crop advisor for further investigation.

Application Instructions

Apply specified rates of **Priaxor® Xemium® brand fungicide** as instructed in **Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions**. **Priaxor** can be applied by ground and aerial application. For best results, thorough coverage of plant materials is required. **Priaxor** can also be applied through sprinkler irrigation equipment, except for use on sugarcane. Application using drip irrigation equipment is permitted in select crops as instructed in **Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions**. Check equipment frequently for calibration. Under low-level disease conditions, the minimum application rates can be used while maximum application rates and shortened spray schedules are recommended for severe or threatening disease conditions.

Cleaning Spray Equipment

Spraying equipment must be cleaned thoroughly before and after applying this product, particularly if a product with potential to injure crops was used prior to **Priaxor**.

For containers 5 gallons or less, shake well prior to use. For containers greater than 5 gallons, recirculate prior to use. Consult BASF Representatives for additional information regarding agitation and recirculation.

Ground Application

Apply **Priaxor** in sufficient water to ensure thorough coverage of foliage, bloom, and fruit. Thorough coverage is required for optimum disease control. Complete coverage of the stem, all the way down to the soil, is required for suppression of soilborne diseases of the stem.

Instructions for Directed or Banded Crop Sprays

The application rates shown in **Table 1. Priaxor® Xemium® brand fungicide Restrictions and Limitations Overview** and **Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions** on this label reflect the amount of product to be applied uniformly over an acre of ground on a broadcast basis. In some crops, **Priaxor** may be used as a directed or banded spray over the rows or plant beds with the alleys or row middles left unsprayed. For such uses, reduce the rate of **Priaxor** in proportion to the area actually sprayed. Make this adjustment to avoid applying the product at use rates higher than permitted on this label.

The following formula may be used to determine the broadcast equivalent rate for doing directed or banded sprays:

sprayed bed width + unsprayed row middle width = total row width

$$\frac{\text{sprayed bed width in inches}}{\text{total row width in inches}} \times \frac{\text{broadcast rate}}{\text{treated acre}} = \frac{\text{band rate}}{\text{field acre}}$$

Example: A directed spray application will be made to 45-inch plant beds that are separated by 15 inches of unsprayed row middles.

45 inches sprayed bed width + 15 inches unsprayed row middles = 60 inches total row width

The calculation to determine the appropriate equivalent rate of product to use for this situation based on a label broadcast rate of 4 fluid ounces product/acre follows:

$$\frac{45 \text{ inches sprayed bed width}}{60 \text{ inches total row width}} \times \frac{4 \text{ fl ozs Priaxor}}{\text{treated acre}} = \frac{3 \text{ fl ozs Priaxor}}{\text{field acre}}$$

Aerial Application

For aerial application in New York State, DO NOT apply within 100 feet of aquatic habitats (such as, but not limited to lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fishponds).

For all crops listed in this label, aerial application can be made and thorough coverage is required to obtain optimum disease control. Avoid applications under conditions when uniform coverage cannot be obtained or when spray drift may occur. **DO NOT** use less than 2 gallons of spray solution per acre. **DO NOT** apply **Priaxor** in spray solutions that are less than 50% water by volume. For aerial applications to sugarcane, **DO NOT** use less than 5 gallons of spray solution per acre. For aerial applications to tree crops, **DO NOT** use less than 10 gallons of spray solution per acre. For all other crops, thorough coverage is required for optimum disease control. The reduced spray volumes used in aerial applications may result in physical incompatibility, reduced disease control, or crop injury from **Priaxor** applications, particularly when tank mixed with other products. Therefore, before making aerial applications test the spray on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application.

Adjuvant or Crop Oil Use Limitations on Corn

Adjuvant crop damage can occur when an adjuvant or crop oil is used after the V8 stage and before the VT stage (the VT stage is defined as when the tassel's last branch is completely visible outside the whorl). If an adjuvant or crop oil is used after the V8 stage and before the VT stage, the grower and user are responsible for contacting the adjuvant source (adjuvant distributor, retailer, or manufacturer) for advice and confirmation that the adjuvant has been tested and proven to be safe for application from V8 to VT corn stage. Refer to adjuvant and/or crop oil labels for specific use directions and restrictions. Always follow the most restrictive label.

Another fungicide or an insecticide may be included in the tank mix if needed and labeled for use on corn. Refer to the tank mix pesticide product labels for specific use

directions and restrictions. Always follow the most restrictive label.

Spray Drift Management

DO NOT spray when conditions favor drift beyond area intended for application. Conditions that may contribute to drift include thermal inversion, wind speed and direction, spray nozzle/pressure combinations, spray droplet size, temperature/humidity, etc. Contact your state extension agent for spray drift prevention guidelines in your area. All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers. Avoiding spray drift at the application site is the responsibility of the applicator.

Aerial Application Methods and Equipment

The interaction of many equipment-related and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

DO NOT apply under circumstances where possible drift to unprotected persons, to food, forage, or other plantings that might be damaged, or crops thereof rendered unfit for sale, use or consumption can occur.

DO NOT release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety or special weather conditions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the fixed wingspan or 90% of rotor blade diameter.
2. Nozzles must always point backward parallel with the airstream and never be pointed downward more than 45 degrees.

Where states have more stringent regulations, they must be observed.

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. Use the largest droplet size consistent with acceptable efficacy. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly or under unfavorable environmental conditions (see **Wind**; **Temperature and Humidity**; and **Temperature Inversions**).

Controlling Droplet Size:

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - **DO NOT** exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is recommended practice unless inconsistent with product efficacy. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid-stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Wind

DO NOT apply at wind speeds greater than 15 mph. Drift potential is lowest when wind speed does not exceed 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Avoid applications below 2 mph due to variable wind direction and high inversion potential. Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

Low humidity and high temperatures increase the evaporation of spray droplets and therefore, the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and/or high temperatures. When making applications in low relative humidity, set up equipment to produce larger droplets in order to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light, variable winds common during inversions.

Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. bodies of water or non-target crops) is minimal and when wind is blowing away from the sensitive areas.

Directions for Use Through Irrigation Systems

Sprayer Preparation

Clean chemical tank and injector system thoroughly. Flush system with clean water.

Application Instructions

Apply **Priaxor® Xemium® brand fungicide** at rates and timings as required in this label.

Use Precautions for Sprinkler Irrigation and Drip Irrigation Applications

- **DO NOT** apply by sprinkler irrigation to sugarcane. For all other crops, this product can be applied through sprinkler irrigation systems including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move irrigation systems equipment. Application using drip irrigation is permitted in select crops as instructed in **Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions**. **DO NOT** apply this product through any other type of irrigation system.
- Add **Priaxor** to the pesticide supply tank containing sufficient water to maintain a continuous flow by the injection equipment. In continuous moving systems, inject this product-water mixture continuously, applying the labeled rate per acre for that crop. **DO NOT** exceed 1/2 inch (13,577 gallons) of water per acre. In stationary or noncontinuous moving systems, inject the product-water mixture in the last 15 to 30 minutes of each set allowing sufficient time for all of the required pesticide to be applied by all the sprinkler heads and applying the labeled rate per acre for that crop. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. Thorough coverage of foliage is required for good control. Maintain good agitation during the entire application period.
- Contact a state extension service specialist, equipment manufacturers or other experts for calibration questions.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

- Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- **DO NOT** connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

Specific Instructions for Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Additives and Tank Mixing Information

Priaxor® Xemium® brand fungicide can be tank mixed with recommended fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives as specified in **Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions**. See berries and small fruits subgroups, cucurbit vegetables group, fruiting vegetables group, grapes, and leafy vegetables group in **Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions** for exceptions.

Under some conditions, the use of additives or adjuvants may improve the performance of **Priaxor**. However, all varieties and cultivars have not been tested with possible tank mix combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, or crop injury may result from mixing **Priaxor** with other products. Therefore, before using any tank mix (fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives), test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application.

When an adjuvant is to be used with this product, BASF recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Consult a BASF representative or local agricultural authorities for more information concerning additives.

If tank mixtures are used, adhere to restrictions due to rates, label instructions and precautions on all labels.

Adjuvant or Crop Oil Use Limitations on Corn

Adjuvant crop damage can occur when an adjuvant or crop oil is used after the V8 stage and before the VT stage (the VT stage is defined as when the tassel's last branch is completely visible outside the whorl). If an adjuvant or crop oil is used after the V8 stage and before the VT stage, the grower and user are responsible for contacting the adjuvant source (adjuvant distributor, retailer, or manufacturer) for advice and confirmation that the adjuvant has been tested and proven to be safe for application from V8 to VT corn stage. Refer to adjuvant and/or crop oil labels for specific use directions and restrictions. Always follow the most restrictive label.

Another fungicide or an insecticide may be included in the tank mix if needed and labeled for use on corn. Refer to the tank mix pesticide product labels for specific use directions and restrictions. Always follow the most restrictive label.

Compatibility Test for Tank Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of label rate per acre:

1. **Water** - For 100 gallons per acre spray volume, use 16 cups (1 gallon) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
2. **Water-dispersible products** (dry flowables, wettable powders, suspension concentrates, or suspoemulsions) - Cap the jar and invert 10 cycles.
3. **Water-soluble products** - Cap the jar and invert 10 cycles.
4. **Emulsifiable concentrates** (oil concentrate or methylated seed oil when applicable) - Cap the jar and invert 10 cycles.
5. **Water-soluble additives** - Cap the jar and invert 10 cycles.
6. Let the solution stand for 15 minutes.
7. **Evaluate** the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. **DO NOT** use any spray solution that could clog spray nozzles.

Mixing Order

1. **Water** - Begin by agitating a thoroughly clean sprayer tank 3/4 full of clean water.
2. **Agitation** - Maintain constant agitation throughout mixing and application.
3. **Inductor** - If an inductor is used, rinse it thoroughly after each component has been added.
4. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. **Water-dispersible products** (such as dry flowables, wettable powders, suspension concentrates including **Priaxor**, or suspo-emulsions) - **For containers 5 gallons or less, shake well prior to use. For containers greater than 5 gallons, recirculate prior to use.** Consult BASF Representatives for additional information regarding agitation and recirculation.
6. **Water-soluble products**
7. **Emulsifiable concentrates** (such as oil concentrates when applicable)
8. **Water-soluble additives** (such as ammonium sulfate [AMS] or urea ammonium nitrate [UAN] when applicable)
9. **Remaining quantity of water**

Make sure that each component is thoroughly mixed and suspended before adding tank mix partners. Maintain constant agitation during application. See **Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions** for more details.

Restrictions and Limitations

- **DO NOT** exceed the maximum product rate (fl ozs/A) per year (season), the maximum rate per application, or the total number of applications of **Priaxor® Xemium® brand fungicide** per year (season) as stated in **Table 1. Priaxor® Xemium® brand fungicide Restrictions and Limitations Overview** and **Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions**. Preharvest interval (PHI) restrictions are also included in these tables.
- **DO NOT** use **Priaxor** in greenhouse or transplant production.
- **For aerial application in New York State, DO NOT apply within 100 feet of aquatic habitats (such as, but not limited to lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fishponds).**
- **Crop Rotation Restriction** - The following crops may be planted immediately following the last application: alfalfa, barley, berries and small fruits, Brassica leafy vegetables, bulb vegetables, corn (all types), cotton, cucurbit vegetables, dried shelled peas and beans, edible-podded legume vegetables, fruiting vegetables (including tomato), grapes, leafy vegetables, mint (spearmint and peppermint), oat, oilseed crops (including flax seed, rapeseed, safflower, and sunflower), peanut, pome fruits, root vegetables, rye, sorghum and millet, soybean, stone fruits, strawberries, succulent shelled peas and beans, sugar beet, sugarcane, tree nuts, tuberous and corm vegetables (including potato), wheat and triticale, and any other crop labeled for direct application of this product.

For rice, **DO NOT** plant sooner than 14 days after the last application.

For all other crops, **DO NOT** plant sooner than 365 days after the last application.

- **Priaxor** is not for sale, distribution, or use in Nassau and Suffolk counties in New York State.

Table 1. Priaxor® Xemium® brand fungicide Restrictions and Limitations Overview*

Crop**	Minimum time from Application to Harvest (PHI) (days)	Maximum Product Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Number of Sequential Applications	Maximum Product Rate per Season (fl ozs/A)
Barley	Apply no later than 50% head emergence (Feekes 10.3, Zadok's 55)	8	2	2	16
Berries and small fruits subgroups	0	8.8	3	2	26.4
Bushberry crop subgroup 13-07B					
Caneberry crop subgroup 13-07A					
Low growing berry (except cranberry) crop subgroup 13-07G					
Small fruit vine climbing (except fuzzy kiwifruit) crop subgroup 13-07F	14				
Brassica leafy vegetables crop subgroups 5A and 5B	3	8.2	3	2	24.6
Bulb vegetables crop group 3-07	7	7.4	3	2	22.2
Corn	21 7 (sweet)	8	2	2	16
Cucurbit vegetables crop group 9	0	8.2	3	1	24.6
Dried shelled peas and beans (except soybeans) crop group 6C	21	8	2	2	16
Edible-podded legume vegetables crop subgroup 6A	7	8	2	2	16
Fruiting vegetables crop group 8	0	8	3	2	24
Grapes	14	6.9	3	2	20.7
Leafy vegetables (except Brassica) crop group 4	1	9.2	3	2	27.6
Oats	Apply no later than 50% head emergence (Feekes 10.3, Zadok's 55)	8	2	2	16
Oilseed crops	21	8	2	2	16

(continued)

Table 1. Priaxor® Xemium® brand fungicide Restrictions and Limitations Overview* (continued)

Crop**	Minimum time from Application to Harvest (PHI) (days)	Maximum Product Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Number of Sequential Applications	Maximum Product Rate per Season (fl ozs/A)
Peanut	14	8	3	2	24
Root vegetables (except sugar beet) crop subgroup 1B	7	8.2	3	2	24.6
Rye	Apply no later than the beginning of flowering (Feekes 10.5, Zadok's 59)	8	2	2	16
Sorghum and millet	21	8	1	1	8
Soybean	21	8	2	2	16
Strawberries	0	8.8	3	2	26.4
Succulent shelled peas and beans	7	8	2	2	16
Sugar beet	7	8	3	2	24
Sugarcane	14	9	2	2	18
Tree nuts crop group 14	14	5.5	3	2	16.5
Tuberous and corm vegetables crop subgroup 1C	7	8	3	2	24
Wheat and triticale	Apply no later than the beginning of flowering (Feekes 10.5, Zadok's 59)	8	2	2	16

* See **Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions** for additional directions.

** For a complete list of crops within a crop group, see **Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions**.

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Barley	Black point (Kernel blight or Head mold) <i>(Cochliobolus sativus,</i> <i>Alternaria spp.)</i> Leaf rust <i>(Puccinia spp.)</i> Net blotch <i>(Pyrenophora teres)</i> Powdery mildew <i>(Blumeria graminis f. sp.</i> <i>hordei)</i> Scald <i>(Rhynchosporium secalis)</i> Septoria leaf and glume blotch <i>(Septoria spp.,</i> <i>Stagonospora spp.)</i> Spot blotch <i>(Cochliobolus sativus)</i> Stem rust <i>(Puccinia graminis f. sp. tritici)</i> Stripe rust <i>(Puccinia striiformis)</i> Tan spot (Yellow leaf spot) <i>(Pyrenophora spp.)</i>	4 to 8*	2	16	Apply no later than 50% head emergence (Feekes 10.3, Zadok's 55)

Application Directions. For optimal disease control, begin applications of **Priaxor** prior to disease development. To maximize yields in cereals, it is important to protect the flag leaf. Apply **Priaxor** immediately after flag leaf emergence for optimum results.

Priaxor does not control Fusarium head blight (head scab) or prevent the reductions in grain quality that can result from this disease. When head blight is a concern, growers should manage this disease with fungicides that are labeled for and effective in managing this disease, and with cultural practices like crop rotation and plowing to reduce crop residues that serve as an inoculum source.

DO NOT harvest barley hay or feed green-chopped barley within 14 days of last application.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 16 fl ozs of **Priaxor** per acre per season. **DO NOT** make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* For early season control of net blotch, Septoria leaf and glume blotch, spot blotch, and tan spot when conditions favor disease development, apply 2 to 4 fl ozs per acre of **Priaxor** either in combination with a herbicide application or when conditions favor disease development. When the 2 to 4 fl ozs early season application rate is used, a second application of **Priaxor** may be required to protect the emerged flag leaf. Environmental conditions for disease or current disease pressure at the time of flag-leaf emergence should be used to determine the **Priaxor** rate for the second application. For high disease pressure, use the higher rate of **Priaxor**.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p>Berries and small fruits subgroups*</p> <p>Bushberry crop subgroup 13-07B Blueberry* (highbush and lowbush) Currant Elderberry Gooseberry Huckleberry Cultivars, varieties, and/or hybrids of these</p> <p>Caneberry crop subgroup 13-07A Blackberry (all varieties) Loganberry Raspberry (black and red) Wild raspberry Cultivars, varieties, and/or hybrids of these</p> <p>Low growing berry (except cranberry) crop subgroup 13-07G Bearberry Bilberry Cloudberry Lingonberry Muntries Partridgeberry Cultivars, varieties, and/or hybrids of these</p>	<p>Alternaria leaf spot and fruit rot (<i>Alternaria</i> spp.)</p> <p>Anthracnose (<i>Colletotrichum</i> spp., <i>Elsinoe</i> spp.)</p> <p>Leaf spot and blotch (<i>Mycosphaerella</i> spp., <i>Septoria</i> spp.)</p> <p>Monilinia blight and mummy berry (<i>Monilinia</i> spp.)</p> <p>Phomopsis leaf spot, twig blight, and fruit rot (<i>Phomopsis</i> spp.)</p> <p>Powdery mildew (<i>Sphaerotheca</i> spp., <i>Microshaera</i> spp., <i>Oidium</i> spp.)</p> <p>Spur blight (<i>Didymella</i> spp., <i>Phoma</i> spp.)</p> <p>Suppression only</p> <p>Botrytis gray mold (<i>Botrytis cinerea</i>)</p> <p>Rust (<i>Pucciniastrum</i> spp., <i>Arthuriomyces</i> spp., <i>Phragmidium</i> spp., <i>Kuehneola</i> spp.)</p>	6 to 8.8	3	26.4	0 (bushberry, caneberry, and low growing berry)
<p>Small fruit vine climbing (except fuzzy kiwifruit) crop subgroup 13-07F Amur river grape Gooseberry Kiwifruit, hardy Maypop Schisandra berry Cultivars, varieties, and/or hybrids of these</p>					14 (small fruit vine climbing)

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Berries and small fruits subgroups* *(continued)*

Application Directions. Begin applications of **Priaxor** prior to onset of disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

DO NOT apply **Priaxor** to any crops in the berries and small fruits subgroups, including blueberries (highbush and low-bush), as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives, or anything other than water.

Mix **Priaxor** with water only for applications to crops listed in the berries and small fruits subgroups.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 26.4 fl ozs of **Priaxor** per acre per season. **DO NOT** apply more than three (3) applications of **Priaxor** per season.

DO NOT make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Brassica leafy vegetables crop subgroups 5A & 5B Head and stem crop subgroup 5A Broccoli Broccoli, Chinese Brussels sprouts Cabbage Cabbage, Chinese Cabbage, Chinese mustard Cauliflower Cavalo broccolo Kohlrabi Leafy greens crop subgroup 5B Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens	Alternaria leaf spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.) Black leg (<i>Phoma lingam</i>) Cercospora leaf spot (<i>Cercospora brassicicola</i>) Powdery mildew (<i>Erysiphe polygoni</i>) Rhizoctonia blight (<i>Rhizoctonia solani</i>) Ring spot (<i>Mycosphaerella brassicicola</i>) White leaf spot (<i>Pseudocercospora capsellae</i>) White rust (<i>Albugo candida</i>) Suppression only Downy mildew (<i>Peronospora parasitica</i>)	6 to 8.2	3	24.6	3

Application Directions. Begin applications of **Priaxor** prior to onset of disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 24.6 fl ozs of **Priaxor** per acre per season. **DO NOT** apply more than three (3) applications of **Priaxor** per season.

DO NOT make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p>Bulb vegetables crop group 3-07*</p> <p>Chive, Chinese, fresh leaves Chive, fresh leaves Daylily, bulb Elegans hosta Fritillaria, bulb Fritillaria, leaves Garlic, bulb Garlic, great-headed, bulb Garlic, serpent, bulb Kurrat Lady's leek Leek Leek, wild Lily, bulb Onion, Beltsville bunching Onion, bulb Onion, Chinese, bulb Onion, fresh Onion, green Onion, macrostem Onion, pearl Onion, potato, bulb Onion, tree, tops Onion, Welsh, tops Shallot, bulb Shallot, fresh leaves Cultivars, varieties, and/or hybrids of these</p>	<p>Powdery mildew (<i>Leveillula taurica</i>) Purple blotch and leaf blight (<i>Alternaria porri</i>) Rust (<i>Puccinia porri</i>) Stemphylium leaf blight and stalk rot (<i>Stemphylium vesicarium</i>) Suppression only Botrytis leaf blight (<i>Botrytis</i> spp.) Botrytis neck rot (<i>Botrytis</i> spp.) Downy mildew (<i>Peronospora destructor</i>)</p>	6 to 7.4	3	22.2	7

Application Directions. Begin applications of **Priaxor** prior to onset of disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 22.2 fl ozs of **Priaxor** per acre per season. **DO NOT** apply more than three (3) applications of **Priaxor** per season.

DO NOT make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Corn Field corn Pop corn Sweet corn Seed production corn	Anthracnose <i>(Colletotrichum graminicola)</i> Eyespot <i>(Kabatiella zea)</i> Gray leaf spot <i>(Cercospora zea-maydis)</i> Northern corn leaf blight <i>(Exserohilum turcicum)</i> Northern corn leaf spot <i>(Cochliobolus carbonum)</i> Physoderma brown spot <i>(Physoderma maydis)</i> Rust, common <i>(Puccinia sorghi)</i> Rust, southern <i>(Puccinia polysora)</i> Southern corn leaf blight <i>(Bipolaris maydis)</i> Yellow leaf blight <i>(Phyllosticta maydis)</i>	4 to 8	2	16	21 (7 sweet corn only)

Application Directions. For optimal disease control, begin applications of **Priaxor** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. For aerial application directions to corn, refer to the **Adjuvant or Crop Oil Use Limitations on Corn** section.

Priaxor may be used with adjuvants. See the **Additives and Tank Mixing Information** and **Mixing Order** sections for more details.

Adjuvant or Crop Oil Use Limitations on Corn. Adjuvant crop damage can occur when an adjuvant or crop oil is used after the V8 stage and before the VT stage (the VT stage is defined as when the tassel's last branch is completely visible outside the whorl). If an adjuvant or crop oil is used after the V8 stage and before the VT stage, the grower and user are responsible for contacting the adjuvant source (adjuvant distributor, retailer, or manufacturer) for advice and confirmation that the adjuvant has been tested and proven to be safe for application from V8 to VT corn stage. Refer to adjuvant and/or crop oil labels for specific use directions and restrictions. Always follow the most restrictive label.

Another fungicide or an insecticide may be included in the tank mix if needed and labeled for use on corn. Refer to the tank mix pesticide product labels for specific use directions and restrictions. Always follow the most restrictive label.

DO NOT harvest for forage within 7 days of last application.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 16 fl ozs of **Priaxor** per acre per season. **DO NOT** make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p>Cucurbit vegetables crop group 9*</p> <p>Chayote Chinese waxgourd Citron melon Cucumber Gherkin Pumpkin Watermelon</p> <p>Edible gourd</p> <p>Chinese okra Cucuzza Hyotan</p> <p>Momordica spp.</p> <p>Balsam apple Balsam pear Bitter melon Chinese cucumber</p> <p>Muskmelon</p> <p>Cantaloupe Casaba Crenshaw melon Golden pershaw melon Honeydew melon Honey balls Mango melon Persian melon Pineapple melon Santaclaus melon Snake melon</p> <p>Summer squash</p> <p>Crookneck squash Scallop squash Straightneck squash Vegetable marrow Zucchini</p> <p>Winter squash</p> <p>Acorn squash Butternut squash Calabaza Hubbard squash Spaghetti squash</p>	<p>Alternaria leaf blight <i>(Alternaria cucumerina)</i></p> <p>Powdery mildew <i>(Podosphaera spp., Sphaerotheca spp., Erysiphe spp.)</i></p> <p>Anthracnose <i>(Colletotrichum orbiculare)</i></p> <p>Cercospora leaf spot <i>(Cercospora citrulina)</i></p> <p>Downy mildew <i>(Pseudoperonospora cubensis)</i></p> <p>Gummy stem blight <i>(Didymella bryoniae)</i></p> <p>Microdochium blight <i>(Plectosporium tabacinum)</i></p> <p>Target leaf spot <i>(Corynespora cassiicola)</i></p>	<p>6 to 8.2</p> <p>8.2</p>	<p>3</p>	<p>24.6</p>	<p>0</p>

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Cucurbit vegetables crop group 9* *(continued)*

Application Directions. For optimal disease control, begin applications of **Priaxor** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

DO NOT apply **Priaxor** to any crops in the cucurbit vegetables group as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives, or anything other than water. See **Resistance Management** section below for tank mix exception for gummy stem blight resistance.

Mix **Priaxor** with water only for applications to crops listed in the cucurbit vegetables group.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 24.6 fl ozs of **Priaxor** per acre per season. **DO NOT** apply more than three (3) applications of **Priaxor** per season.

DO NOT make more than one (1) application of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide. In areas where gummy stem blight resistance to **Group 7** or **Group 11** fungicides have been confirmed, tank mix with chlorothalonil at full label rates and adhering to all label precautions.

* Not registered for use in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Dried shelled peas and beans (except soybeans) crop group 6C* Broad bean Chickpea Guar Lablab bean Lentil Pigeon pea <i>Lupinus</i> spp. Grain lupin Sweet lupin White lupin <i>Phaseolus</i> spp. Field bean Kidney bean Lima bean Navy bean Pink bean Pinto bean Tepary bean <i>Pisum</i> spp. Field pea	Anthracnose (<i>Colletotrichum</i> spp.) Alternaria leaf and pod spot (<i>Alternaria</i> spp.) Ascochyta blight (<i>Phoma exigua</i> , <i>Ascochyta</i> spp.) Asian soybean rust (<i>Phakopsora pachyrhizi</i>) Botrytis gray mold (<i>Botrytis cinerea</i>) Cercospora leaf spot (<i>Cercospora</i> spp.) Downy mildew (<i>Phytophthora nicotianae</i>) Mycosphaerella blight (<i>Mycosphaerella</i> spp.) Powdery mildew (<i>Erysiphe polygoni</i>) Rust (<i>Uromyces appendiculatus</i>)	4 to 8	2	16	21
<i>Vigna</i> spp. Adzuki bean Blackeyed pea Catjang Cowpea Crowder pea Moth bean Mung bean Rice bean Southern pea Urd bean	Suppression only White mold (<i>Sclerotinia sclerotiorum</i>)	6 to 8			

Application Directions. For optimal disease control, begin applications of **Priaxor** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

Bean forage, bean hay, pea vines, and pea hay may be fed no sooner than 14 days after last application.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Dried shelled peas and beans (except soybeans) crop group 6C* *(continued)*

Use of Adjuvants and Other Products as Mixes with Priaxor.

The use of adjuvants or additives may improve the performance of **Priaxor** on dried shelled peas and beans. However, under certain conditions, mixtures of **Priaxor** with adjuvants, additives and/or other products may cause crop injury.

DO NOT use **Priaxor** with:

- Emulsifiable concentrate (EC) or solvent-based formulation products.
- Crop oil concentrate (COC), methylated seed oil (MSO), organosilicone (OS), MSO/OS blend adjuvants.
- Nonionic surfactant (NIS) adjuvant products that acidify or enhance plant penetration on dried shelled peas and beans.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, crop injury, or incompatibility due to additives, adjuvants or other products used in combination with **Priaxor** may result from mixing **Priaxor** with other products. Refer also to the **Conditions of Sale and Warranty** section of this label.

To minimize the likelihood of crop injury, BASF recommends testing **Priaxor** in combination with other products for crop safety on a small portion of the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult a BASF representative for more information concerning additives or adjuvants.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 16 fl ozs of **Priaxor** per acre per season. **DO NOT** make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use on dried shelled peas and beans in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Edible-podded legume vegetables crop subgroup 6A* Jack bean Pigeon pea Soybean (immature seed) Sword bean <i>Phaseolus spp.</i> Runner bean Snap bean Wax bean <i>Pisum spp.</i> Dwarf pea Edible-podded pea Snowpea Sugar snap pea <i>Vigna spp.</i> Asparagus bean Chinese longbean Moth bean Yardlong bean	Anthracnose (<i>Colletotrichum spp.</i>) Alternaria leaf and pod spot (<i>Alternaria spp.</i>) Ascochyta blight (<i>Phoma exigua</i> , <i>Ascochyta spp.</i>) Asian soybean rust (<i>Phakopsora pachyrhizi</i>) Botrytis gray mold (<i>Botrytis cinerea</i>) Cercospora leaf spot (<i>Cercospora spp.</i>) Downy mildew (<i>Phytophthora nicotianae</i>) Mycosphaerella blight (<i>Mycosphaerella spp.</i>) Powdery mildew (<i>Erysiphe polygoni</i>) Rust (<i>Uromyces appendiculatus</i>)	4 to 8	2	16	7
	Suppression only White mold (<i>Sclerotinia sclerotiorum</i>)	6 to 8			
<p>Application Directions. For optimal disease control, begin applications of Priaxor prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.</p> <p>Bean forage, bean hay, pea vines, and pea hay may be fed no sooner than 14 days after last application.</p>					

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Edible-podded legume vegetables crop subgroup 6A* *(continued)*

Use of Adjuvants and Other Products as Mixes with Priaxor.

The use of adjuvants or additives may improve the performance of **Priaxor** on edible podded legumes. However, under certain conditions, mixtures of **Priaxor** with adjuvants, additives and/or other products may cause crop injury.

DO NOT use **Priaxor** with:

- Emulsifiable concentrate (EC) or solvent-based formulation products.
- Crop oil concentrate (COC), methylated seed oil (MSO), organosilicone (OS), MSO/OS blend adjuvants.
- Nonionic surfactant (NIS) adjuvant products that acidify or enhance plant penetration on edible podded legumes.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, crop injury, or incompatibility due to additives, adjuvants or other products used in combination with **Priaxor** may result from mixing **Priaxor** with other products. Refer also to the **Conditions of Sale and Warranty** section of this label.

To minimize the likelihood of crop injury, BASF recommends testing **Priaxor** in combination with other products for crop safety on a small portion of the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult a BASF representative for more information concerning additives or adjuvants.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 16 fl ozs of **Priaxor** per acre per season. **DO NOT** make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Fruiting vegetables crop group 8 Eggplant Ground cherry Pepino Pepper (all varieties) Tomatillo Tomato	Anthracnose (<i>Colletotrichum coccodes</i>) Black mold (<i>Alternaria alternata</i>) Early blight (<i>Alternaria solani</i>) Septoria leaf spot (<i>Septoria lycopersici</i>) Target spot (<i>Corynespora cassiicola</i>)	4 to 8 or 4 to 8 fl ozs per 100 gallons of spray volume (dilute)*	3	24	0
	Powdery mildew (<i>Leveillula taurica</i>)	6 to 8			
	Suppression only	4 to 8			
	Botrytis gray mold (<i>Botrytis cinerea</i>) Rhizoctonia stem rot** (<i>Rhizoctonia solani</i>) Sclerotinia stem rot, White mold (<i>Sclerotinia sclerotiorum</i>) Southern blight** (<i>Sclerotium rolfsii</i>)	4 to 8 or 4 to 8 fl ozs per 100 gallons of spray volume (dilute)*			
	Suppression only	8			
	Late blight (<i>Phytophthora infestans</i>)				

Application Directions. For optimal disease control, begin applications of **Priaxor** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

Applications using drip irrigation systems may provide disease suppression. The level and consistency of suppression from drip line applications varies with the soil type, level of inoculum, irrigation volumes, environment and other factors. See your local BASF representative for details on drip irrigation use in your area.

* For applications based on dilute volume, plants should be sprayed to runoff. Apply a minimum of 20 gallons of spray volume per acre, and increase the spray volume as the plants grow during the season. Spray volume should be proportional to the amount of plant tissue to be covered such that 100 gallons of spray per acre is used on mature plants.

Use of Adjuvants and Other Products as Mixes with Priaxor.

Priaxor can be used with nonionic surfactants at their lowest label rate up to 0.125%. When **Priaxor** is mixed with buffering agents and foliar nutrients, the pH of the final spray solution must be greater than 5.5.

DO NOT mix **Priaxor** with the following products:

- Emulsifiable concentrate (EC) formulation or solvent-based formulation products.
- Crop oil concentrate (COC), methylated seed oil (MSO), organosilicone (OS) or MSO/OS blended adjuvant products.

For **Priaxor** applications to **fresh market tomatoes** at less than 20 gallons per acre, **DO NOT** mix **Priaxor** with any other products, adjuvants, additives, nutrients or anything other than water.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Fruiting vegetables crop group 8 *(continued)*

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, crop injury, or incompatibility due to additives, adjuvants or other products used in combination with **Priaxor** may result from mixing **Priaxor** with other products. Refer also to the **Conditions of Sale and Warranty** section of this label.

To minimize the likelihood of crop injury, BASF recommends testing **Priaxor** in combination with other products for crop safety on a small portion of the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult a BASF representative for more information concerning additives or adjuvants.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 24 fl ozs of **Priaxor** per acre per season. **DO NOT** apply more than three (3) applications of **Priaxor** per season.

DO NOT make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

** Not registered for use in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Grapes* (except Concord and Noiret) See Crop Safety Precaution below	Powdery mildew <i>(Erysiphe necator)</i>	4 to 6	3	18	14
	Angular leaf spot <i>(Mycosphaerella angulata)</i> Anthracnose <i>(Elsinoe ampelina)</i> Black rot <i>(Guignardia bidwellii)</i> Downy mildew <i>(Plasmopara viticola)</i> Leaf blight <i>(Pseudocercospora vitis)</i> Ripe rot <i>(Colletotrichum gloeosporioides)</i> Suppression only Botrytis gray mold <i>(Botrytis cinerea)</i> Phomopsis cane and Leaf spot <i>(Phomopsis viticola)</i> Aids in control: Summer bunch rot <i>(Aspergillus spp., Alternaria spp., Botrytis spp., Cladosporium spp, Penicillium spp., Rhizopus spp.)</i>	6 to 6.9	3	20.7	

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Grapes* (except Concord and Noiret) *(continued)*

Application Directions. For powdery mildew control, begin applications of **Priaxor** as of bud break or prior to onset of disease. When using the lower rate, continue on a 14 to 21 day interval. Longer spray intervals for powdery mildew control may be possible with higher rates of **Priaxor**. The effectiveness of longer spray intervals will depend on the current powdery mildew infection level in the field, the amount of disease pressure after application and factors such as crop growth stage and rate of growth.

For black rot and downy mildew control, begin applications of **Priaxor** before pre-bloom and continue on a 10 to 14 day interval.

Begin applications targeted against **Botrytis gray mold** prior to disease development when conditions favor the development of Botrytis such as during early bloom, bunch preclosure and veraison. Use the higher rates from bunch preclosure to veraison. Use shorter intervals when disease pressure is high.

For all other diseases listed, begin applications of **Priaxor** prior to onset of disease development and continue on a 10 to 14 day interval. Use the higher rate and the shorter interval when disease pressure is high.

DO NOT apply **Priaxor** to grapes as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives, or anything other than water.

Mix **Priaxor** with water only for applications to grapes.

Crop Safety Precaution. DO NOT use Priaxor on Concord, Noiret and NY73.0136.17 due to possible foliar injury. Possible foliar injury could occur to Worden, Fredonia, Niagara, Steuben, Rougeon or related varieties. Not all varieties have been thoroughly tested.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 20.7 fl ozs of **Priaxor** per acre per season. **DO NOT** make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Leafy vegetables (except Brassica) crop group 4* Amaranth Arugula Cardoon Celery Celery, Chinese Celtuce Chervil Chrysanthemum (edible-leaved and garland) Corn salad Cress (garden and Upland) Dandelion Dock Endive Fennel, Florence Lettuce (head and leaf) Orach Parsley Purslane (garden and winter)	Alternaria leaf spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.) Ascochyta leaf spot (<i>Ascochyta</i> spp.) Cercospora leaf spot (<i>Cercospora</i> spp.) Downy mildew (<i>Peronospora</i> spp.) Phoma (<i>Phoma</i> spp.) Powdery mildew (<i>Erysiphe</i> spp., <i>Phyllactinia</i> spp., <i>Sphaerotheca</i> spp.) Rust (<i>Puccinia</i> spp., <i>Uromyces</i> spp.) Septoria leaf spot (<i>Septoria</i> spp.) White rust (<i>Albugo</i> spp.)	6 to 9.2	3	27.6	1
Radicchio (red chicory) Rhubarb Spinach Spinach (New Zealand and vine) Swiss chard	Suppression only Botrytis rot (<i>Botrytis</i> spp.) Lettuce downy mildew (<i>Bremia</i> spp.) Lettuce drop caused by <i>Sclerotinia minor</i>	9.2			

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Leafy vegetables (except Brassica) crop group 4* *(continued)*

Application Directions. Begin applications of **Priaxor** prior to onset of disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

Tank Mix Restrictions

Spinach (all varieties). **DO NOT** apply **Priaxor** to spinach as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives, or anything other than water.

Mix **Priaxor** with water only for applications to spinach (all varieties).

Leafy vegetables (except spinach). It is impossible for BASF to test all varieties of leafy vegetables for sensitivity to **Priaxor** under all environments and all potential product mixture combinations.

It is impossible for BASF to test all varieties of leafy vegetables for sensitivity to **Priaxor** under all environments and all potential product mixture combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Proceed with caution with regard to **Priaxor** use, particularly in tank mixes and/or adjuvant combinations on leafy vegetables. To reduce the risk of leafy vegetable injury, BASF recommends testing **Priaxor** or **Priaxor** tank mixtures on a small portion of the crop before broadscale use.

To the extent consistent with applicable law, the user assumes all risks associated with adding products to the **Priaxor** spray solution. Refer also to the **Conditions of Sale and Warranty** section of this label.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 27.6 fl ozs of **Priaxor** per acre per season. **DO NOT** apply more than three (3) applications of **Priaxor** per season.

DO NOT make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Oats*	Crown rust <i>(Puccinia coronata)</i> Helminthosporium leaf spot <i>(Dreschlera avenae)</i> Leaf blotch <i>(Pyrenophora avenae)</i> Leaf rust <i>(Puccinia spp.)</i> Septoria blotch and stem rot <i>(Septoria spp., Phaeosphaeria spp., Stagonospora spp.)</i> Spot blotch <i>(Bipolaris spp.)</i> Stem rust <i>(Puccinia graminis f. sp. avenae)</i>	4 to 8**	2	16	Apply no later than 50% head emergence (Feekes 10.3, Zadok's 55)

Application Directions. For optimal disease control, begin applications of **Priaxor** prior to disease development. To maximize yields in cereals, it is important to protect the flag leaf. Apply **Priaxor** immediately after flag leaf emergence for optimum results.

Priaxor does not control Fusarium head blight (head scab) or prevent the reductions in grain quality that can result from this disease. When head blight is a concern, growers should manage this disease with fungicides that are labeled for and effective in managing this disease, and with cultural practices like crop rotation and plowing to reduce crop residues that serve as an inoculum source.

DO NOT harvest oat hay or feed green-chopped oats within 14 days of last application.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 16 fl ozs of **Priaxor** per acre per season. **DO NOT** make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

** For early season control of leaf blotch, Septoria blotch and stem rot, and spot blotch when conditions favor disease development, apply 2 to 4 fl ozs per acre of **Priaxor** either in combination with a herbicide application or when conditions favor disease development. When the 2 to 4 fl ozs early season application rate is used, a second application of **Priaxor** may be required to protect the emerged flag leaf. Environmental conditions for disease or current disease pressure at the time of flag-leaf emergence should be used to determine the **Priaxor** rate for the second application. For high disease pressure, use the higher rate of **Priaxor**.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Oilseed crops Flax seed	Pasm <i>(Septoria linicola)</i>	4 to 8	2	16	21
Rapeseed (cultivars, varieties, and/or hybrids, including canola and crambe)	Blackleg <i>(Leptosphaeria maculans)</i> Blackspot <i>(Alternaria spp)</i> Suppression only White mold/ Sclerotinia stem rot <i>(Sclerotinia sclerotiorum)</i>				
Safflower	<i>Alternaria spp.</i> <i>Septoria spp.</i> Suppression only <i>Sclerotinia spp.</i>				
Sunflower	Alternaria leaf spot <i>(Alternaria spp.)</i> Cercospora leaf spot <i>(Cercospora helianthi)</i> Powdery mildew <i>(Erysiphe cichoracearum)</i> Rust <i>(Puccinia helianthi, Uromyces spp.)</i> Septoria leaf spot <i>(Septoria spp.)</i> White rust <i>(Albugo tragopogonis)</i> Suppression only Sclerotinia head blight <i>(Sclerotinia sclerotiorum)</i>				

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Oilseed crops Borage Calendula Castor oil plant Chinese tallowtree Cottonseed* Crambe Cuphea Echium Euphorbia Evening primrose Gold of pleasure (camelina) Hare's ear mustard Jojoba Lesquerella Lunaria Meadowfoam Milkweed Mustard seed Niger seed Oil radish Poppy seed Rose hip Sesame Stokes aster Sweet rocket Tallowwood Tea oil plant Vernonia	<i>Alternaria</i> spp. <i>Septoria</i> spp. Suppression only <i>Sclerotinia</i> spp.	4 to 8	2	16	21

Application Directions for Rapeseed. For the control of blackleg, apply **Priaxor** at the 2 to 4 leaf stage. For optimal control of blackspot, apply **Priaxor** at early pod development. For suppression of Sclerotinia, apply **Priaxor** at 20% to 50% flowering or prior to the onset of disease. A second application may be made 14 days later if weather conditions are favorable for disease development.

Application Directions for Other Oilseed Crops. Begin applications of **Priaxor** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

No livestock feeding restrictions.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 16 fl ozs of **Priaxor** per acre per season. **DO NOT** make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Peanut	Early leaf spot <i>(Cercospora arachidicola)</i> Late leaf spot <i>(Cercosporidium personatum)</i> Pepper spot <i>(Leptospherulina crassiasca)</i> Rust <i>(Puccinia arachidis)</i> Web blotch <i>(Phoma arachidicola)</i>	4 to 8	3	24	14
	Rhizoctonia limb rot, Peg rot and Pod rot <i>(Rhizoctonia solani)</i> Sclerotium rot - Southern stem rot, Southern blight and White mold <i>(Sclerotium rolfsii)</i> Suppression only Sclerotinia blight <i>(Sclerotinia minor)</i> Cylindrocladium black rot <i>(Cylindrocladium crotalariae)</i>	8			

Application Directions. For control of early and late leaf spot, pepper spot, rust and web blotch, begin applications of **Priaxor** prior to disease development and continue on a 14 to 21 day interval.

For control of Rhizoctonia and Sclerotium, begin applications of **Priaxor** prior to disease development and continue on a 14 to 28 day interval.

Use the higher rate and/or shorter spray interval when disease pressure is high or in fields with a history of disease.

Priaxor use in mixes with oil, methylated seed oil, crop oil concentrate and/or silicone-containing adjuvants may cause crop injury under certain conditions.

Peanut meal may be fed. **DO NOT** graze or harvest for forage use.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 24 fl ozs of **Priaxor** per acre per season. **DO NOT** apply more than three (3) applications of **Priaxor** per season.

DO NOT make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Root vegetables (except sugar beet) crop subgroup 1B* Beet, garden Burdock, edible Carrot Celeriac Chervil, turnip-rooted Chicory Ginseng Horseradish Parsley, turnip-rooted Parsnip Radish Radish, Oriental Rutabaga Salsify Salsify, black Salsify, Spanish Skirret Turnip	Alternaria leaf spot/blight <i>(Alternaria spp.)</i> Powdery mildew <i>(Erysiphe spp., Leveillula spp.)</i>	6 to 8.2	3	24.6	7
	Cercospora leaf spot/blight <i>(Cercospora spp.)</i> Suppression only Sclerotinia white mold/ cottony rot <i>(Sclerotinia sclerotiorum)</i> Southern blight <i>(Sclerotium rolfsii)</i>	8.2			

Application Directions. For optimal disease control, begin applications of **Priaxor** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 24.6 fl ozs of **Priaxor** per acre per season. **DO NOT** apply more than three (3) applications of **Priaxor** per season.

DO NOT make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Rye	Black point (Kernel blight or Head mold) <i>(Cochliobolus sativus,</i> <i>Alternaria spp.)</i> Leaf rust <i>(Puccinia spp.)</i> Net blotch <i>(Pyrenophora teres)</i> Powdery mildew <i>(Blumeria graminis f. sp. secalis)</i> Scald <i>(Rhynchosporium secalis)</i> Septoria leaf and glume blotch <i>(Septoria spp.,</i> <i>Stagonospora spp.)</i> Spot blotch <i>(Cochliobolus sativus)</i> Stem rust <i>(Puccinia graminis f. sp. secalis)</i> Stripe rust <i>(Puccinia striiformis)</i> Tan spot (Yellow leaf spot) <i>(Pyrenophora spp.)</i>	4 to 8*	2	16	Apply no later than beginning of flowering (Feekes 10.5, Zadok's 59)

Application Directions. For optimal disease control, begin applications of **Priaxor** prior to disease development. To maximize yields in cereals, it is important to protect the flag leaf. Apply **Priaxor** immediately after flag leaf emergence for optimum results.

Priaxor does not control Fusarium head blight (head scab) or prevent the reductions in grain quality that can result from this disease. When head blight is a concern, growers should manage this disease with fungicides that are labeled for and effective in managing this disease, and with cultural practices like crop rotation and plowing to reduce crop residues that serve as an inoculum source.

DO NOT harvest rye hay or feed green-chopped rye within 7 days of last application.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 16 fl ozs of **Priaxor** per acre per season. **DO NOT** make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* For early season control of net blotch, Septoria leaf and glume blotch, spot blotch, and tan spot when conditions favor disease development, apply 2 to 4 fl ozs per acre of **Priaxor** either in combination with a herbicide application or when conditions favor disease development. When the 2 to 4 fl ozs early season application rate is used, a second application of **Priaxor** may be required to protect the emerged flag leaf. Environmental conditions for disease or current disease pressure at the time of flag-leaf emergence should be used to determine the **Priaxor** rate for the second application. For high disease pressure, use the higher rate of **Priaxor**.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Sorghum* Milo* Millet* (pearl and proso)	Anthracnose <i>(Colletotrichum graminicola)</i> Gray leaf spot and Cercospora leaf spot <i>(Cercospora spp.)</i> Northern leaf blight <i>(Exserohilum turcicum)</i> Rust <i>(Puccinia spp.)</i> Southern leaf blight and Bipolaris leaf spot <i>(Bipolaris spp.)</i>	4 to 8	1	8	21
<p>Application Directions. For optimal disease control, begin applications of Priaxor prior to disease development. Use the higher rate when disease pressure is high.</p> <p>Resistance Management. To limit the potential for development of resistance, DO NOT apply more than 8 fl ozs of Priaxor per acre per season. DO NOT apply more than one (1) application of Priaxor per season.</p> <p>* Not registered for use in California.</p>					

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Soybean*	Alternaria leaf spot <i>(Alternaria spp.)</i>	4 to 8	2	16	21
	Anthracnose <i>(Colletotrichum truncatum)</i>				
	Asian soybean rust <i>(Phakopsora pachyrhizi)</i>				
	Brown spot <i>(Septoria glycines)</i>				
	Cercospora blight <i>(Cercospora kikuchii)</i>				
	Frogeye leaf spot <i>(Cercospora sojina)</i>				
	Pod and stem blight <i>(Diaporthe phaseolorum)</i>				
	Rhizoctonia aerial blight <i>(Rhizoctonia solani)</i>				
	Target Spot <i>(Corynespora cassicola)</i>				
	Suppression only				
Sclerotinia blight (white mold) <i>(Sclerotinia sclerotiorum)</i>	8				
Suppression only					
Southern blight <i>(Sclerotium rolfsii)</i>					

Application Directions. For optimal disease control, begin applications of **Priaxor** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

For adequate control of soybean rust, apply **Priaxor** prior to infection.

Priaxor may be used with adjuvants. See the **Additives and Tank Mixing Information** and **Mixing Order** sections for more details.

Soybean forage may be fed no sooner than 14 days after last application. Soybean hay may be fed no sooner than 21 days after last treatment.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 16 fl ozs of **Priaxor** per acre per season. **DO NOT** make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Strawberries*	Leaf spot <i>(Mycosphaerella fragariae, Ramularia tulasnei)</i> Powdery mildew <i>(Sphaerotheca macularis)</i>	6 to 8.8	3	26.4	0
	Anthracnose <i>(Colletotrichum spp.)</i> <u>Suppression only</u> Botrytis gray mold <i>(Botrytis cinerea)</i>	8.8			

Application Directions. Begin applications of **Priaxor** no later than 10% bloom, or prior to disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

The restricted entry interval (REI) for treated strawberries is **12 hours**. Refer to the **Agricultural Use Requirements** section for PPE required for early entry to treated areas as permitted under the Worker Protection Standard.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 26.4 fl ozs of **Priaxor** per acre per season. **DO NOT** apply more than three (3) applications of **Priaxor** per season.

DO NOT make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Succulent shelled peas and beans* Pigeon pea <i>Phaseolus spp.</i> Lima bean, green <i>Pisum spp.</i> English pea Garden pea Green pea Broad bean <i>Vigna spp.</i> Blackeyed pea Cowpea Southern pea	Anthracnose (<i>Colletotrichum spp.</i>)	4 to 8	2	16	7
	Alternaria leaf and pod spot (<i>Alternaria spp.</i>)				
	Ascochyta blight (<i>Phoma exigua</i> , <i>Ascochyta spp.</i>)				
	Asian soybean rust (<i>Phakopsora pachyrhizi</i>)				
	Cercospora leaf spot (<i>Cercospora spp.</i>)				
	Downy mildew (<i>Phytophthora nicotianae</i>)				
	Mycosphaerella blight (<i>Mycosphaerella spp.</i>)				
	Powdery mildew (<i>Erysiphe polygoni</i>)				
	Rust (<i>Uromyces appendiculatus</i>)				
	Suppression only Botrytis gray mold (<i>Botrytis cinerea</i>)		4 to 8		
Suppression only White mold (<i>Sclerotinia sclerotiorum</i>)		6 to 8			

Application Directions. For optimal disease control, begin applications of **Priaxor** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

Bean forage, bean hay, pea vines, and pea hay may be fed no sooner than 14 days after last application.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Succulent shelled peas and beans* *(continued)*

Use of Adjuvants and Other Products as Mixes with Priaxor.

The use of adjuvants or additives may improve the performance of **Priaxor** on succulent shelled peas and beans. However, under certain conditions, mixtures of **Priaxor** with adjuvants, additives and/or other products may cause crop injury.

DO NOT use **Priaxor** with:

- Emulsifiable concentrate (EC) or solvent-based formulation products.
- Crop oil concentrate (COC), methylated seed oil (MSO), organosilicone (OS), MSO/OS blend adjuvants.
- Nonionic surfactant (NIS) adjuvant products that acidify or enhance plant penetration on succulent shelled peas and beans.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, crop injury, or incompatibility due to additives, adjuvants or other products used in combination with **Priaxor** may result from mixing **Priaxor** with other products. Refer also to the **Conditions of Sale and Warranty** section of this label.

To minimize the likelihood of crop injury, BASF recommends testing **Priaxor** in combination with other products for crop safety on a small portion of the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult a BASF representative for more information concerning additives or adjuvants.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 16 fl ozs of **Priaxor** per acre per season. **DO NOT** make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Sugar beet (leaves, roots and tops)	Cercospora leaf spot (<i>Cercospora beticola</i>) Powdery mildew (<i>Erysiphe betae</i>) Rhizoctonia stem canker and crown rot* (<i>Rhizoctonia solani</i>)	6 to 8*	3	24	7

Application Directions. For optimal disease control, begin applications of **Priaxor** prior to disease development and continue on a 14 day interval. Use the higher rate when disease pressure is high.

Sugar beet leaves, roots and tops may be fed no sooner than 7 days after last application.

Use of Adjuvants and Other Products as Mixes with Priaxor.

The use of adjuvants or additives may improve the performance of **Priaxor** on sugar beets. However, under certain conditions, mixtures of **Priaxor** with adjuvants, additives and/or other products may cause crop injury.

DO NOT use **Priaxor** with:

- Emulsifiable concentrate (EC) or solvent-based formulation products.
- Crop oil concentrate (COC), methylated seed oil (MSO), organosilicone (OS), MSO/OS blend adjuvants.
- Nonionic surfactant (NIS) adjuvant products that acidify or enhance plant penetration on sugar beets.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility due to additives, adjuvants or other products used in combination with **Priaxor** may result from mixing **Priaxor** with other products. Refer also to the **Conditions of Sale and Warranty** section of this label.

To minimize the likelihood of crop injury, BASF recommends testing **Priaxor** in combination with other products for crop safety on a small portion of the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult a BASF representative for more information concerning additives or adjuvants.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 24 fl ozs of **Priaxor** per acre per season. **DO NOT** make more than one (1) application of **Priaxor** before the 4-leaf stage of plant growth. After the 4-leaf stage of plant growth, **DO NOT** make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

*Rhizoctonia stem canker and crown rot are registered for use in California at 8 fl ozs/A.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Sugarcane*	Brown rust <i>(Puccinia melanocephala)</i> Orange rust <i>(Puccinia kuehnii)</i>	4 to 9	2	18	14

Application Directions. Begin applications of **Priaxor** at the first sign of disease. Repeat applications on 14 to 28 day intervals as needed if conditions for rust infection continue. Use the shorter interval and/or the higher rate when disease pressure is high.

DO NOT apply this product through any type of irrigation system.

DO NOT feed sugarcane commodities treated with **Priaxor** to livestock.

Priaxor can be applied by ground or air. When applying by air, **DO NOT** use less than 5 gallons of spray solution per acre.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 18.0 fl ozs of **Priaxor** per acre per season. **DO NOT** apply more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p>Tree nuts crop group 14*</p> <p>Almond Beechnut Brazil nut Butternut Cashew Chestnut Chinquapin Filbert Hickory nut Macadamia nut Pecan Pistachio Walnut, black Walnut, English</p>	<p>Alternaria late blight (<i>Alternaria</i> spp.)</p> <p>Anthracnose (<i>Colletotrichum</i> spp.)</p> <p>Leaf rust (<i>Tranzschelia discolor</i>)</p> <p>Scab (<i>Cladosporium carpophilum</i>, <i>C. caryigenum</i>)</p> <p>Shothole (<i>Wilsonomyces carpophilus</i>)</p> <p>Suppression only</p> <p>Botrytis blossom and shoot blight (<i>Botrytis cinerea</i>)</p> <p>Brown rot/Blossom Blight (<i>Monilinia</i> spp.)</p> <p>Green fruit rot/Jacket rot (<i>Botrytis cinerea</i>, <i>Sclerotinia sclerotiorum</i>, <i>Monilinia laxa</i>)</p> <p>Hull rot (<i>Rhizopus stolonifer</i> and <i>Monilinia</i> spp.)</p> <p>Panicle and shoot blight (<i>Botryosphaeria dothidea</i>)</p>	5.5	3	16.5	14

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Tree nuts crop group 14* *(continued)*

Application Directions. For almond, begin applications of **Priaxor** prior to the onset of disease development, and continue on a 7 to 14 day interval up to 14 days before harvest.

For filbert, begin applications at budswell to budbreak, or prior to infection and onset of disease development. Continue on a 7 to 14 day interval to cover and protect new growth.

For pecan, begin applications of **Priaxor** prior to onset of disease development and continue on a 7 to 21 day interval for the control of scab. Use the shorter interval and/or the higher rate when disease pressure is high.

For pistachio, apply **Priaxor** prior to onset of disease development and continue on a 10 to 30 day interval. Use the higher rate and shorter intervals when disease pressure is high.

For all other crops listed, apply **Priaxor** prior to disease development and continue on a 7 to 28 day interval. In all cases use the shorter interval when shoot growth is rapid.

No restriction on livestock feeding of almond hulls.

For aerial application to tree nuts, **DO NOT** use less than 10 gallons of spray solution per acre.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 16.5 fl ozs of **Priaxor** per acre per season. **DO NOT** apply more than three (3) applications of **Priaxor** per season.

DO NOT make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Tuberous and corm vegetables crop subgroup 1C Arracacha Arrowroot Cassava (bitter and sweet) Chayote (root) Chinese artichoke Chufa Dasheen (taro) Edible canna Ginger Jerusalem artichoke Leren	Black dot (<i>Colletotrichum coccodes</i>) Brown spot and Black pit** (<i>Alternaria alternata</i>) Early blight (<i>Alternaria solani</i>) Suppression only Botrytis gray mold** (<i>Botrytis cinerea</i>) Late blight** (<i>Phytophthora infestans</i>)	4 to 8	3	24*	7
Potato Sweet potato Tanier True yam Turmeric Yam bean	Leaf spot (<i>Cercospora</i> spp.) Powdery mildew (<i>Erysiphe</i> spp., <i>Leveillula taurica</i>) Rust (<i>Uromyces</i> spp., <i>Puccinia</i> spp.) Suppression only White mold (<i>Sclerotinia sclerotiorum</i>)	6 to 8			

Application Directions. For optimal disease control, begin applications of **Priaxor** prior to disease development and continue on a 7 to 14 day interval. The lower rate and longer interval can be used early season prior to the observance of symptoms and when disease pressure is low.

Use the higher rates and shorter intervals once disease has been confirmed in your area or weather conditions are conducive to disease development.

Use of Adjuvants and Other Products as Mixes with Priaxor. The use of additives or adjuvants may improve the performance of **Priaxor** on tuberous and corm vegetables.

However, BASF evaluations also indicate that under some conditions, applications of **Priaxor** in combination with certain rates of organosilicone-based or oil-containing (petroleum, crop or methylated seed oil) additives or adjuvants, can cause crop injury. Crop injury also may result from applications of **Priaxor** mixed with other products that have solvent-based formulations that increase penetration.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, crop injury, or incompatibility due to additives, adjuvants or other products used in combination with **Priaxor** may result from mixing **Priaxor** with other products. Refer also to the **Conditions of Sale and Warranty** section of this label.

To minimize the likelihood of crop injury, BASF recommends testing **Priaxor** in combination with other products for crop safety on a small portion of the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult a BASF representative for more information concerning additives or adjuvants.

* The maximum product rate per season includes the combination of infurrow and foliar uses (for above-listed crops, infurrow use is permitted in potato only).

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Tuberous and corm vegetables crop subgroup 1C *(continued)*

No livestock feeding restrictions.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 24 fl ozs of **Priaxor** per acre per season. **DO NOT** apply more than three (3) applications of **Priaxor** per season.

DO NOT make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

Instructions for Infurrow Use to Aid in the Control of Soilborne Rhizoctonia in Potatoes

Use 0.6 fl oz of **Priaxor** per 1000 row feet (for applications on 32-inch or 34-inch rows, the maximum application rate is 0.48 fl oz/1000 row feet). Refer to the chart below to determine the rate per acre. Apply at planting as an infurrow spray by directing spray pattern to uniformly cover seed pieces and surrounding soil. The spray pattern should be a 4 to 8 inch band that is applied to the seed piece prior to being covered with soil.

When Rhizoctonia disease pressure conditions are expected to be severe or if the field has a history of Rhizoctonia infestations, use **Priaxor** at 0.48 to 0.6 fl oz product per 1000 row feet and/or tank mix with a fungicide having a different mode of action.

Use a minimum volume of application of 5 gallons of water per acre.

** Not registered for use in California.

Product Rate per 1000 row feet (fl oz product)	Product Rate per Acre (fl ozs product)			
	34-inch rows or less	36-inch rows	38-inch rows	40-inch rows
0.6	See footnote ¹	8.1	7.7	7.3

¹ For 34-inch rows or less, use a maximum of 0.48 fl oz product per 1000 row feet.

(continued)

Table 2. Priaxor® Xemium® brand fungicide Crop-specific Directions *(continued)*

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Wheat and triticale	Black point (Kernel blight or Head mold) (<i>Cochliobolus sativus</i> , <i>Alternaria</i> spp.) Leaf rust (<i>Puccinia</i> spp.) Powdery mildew (<i>Blumeria graminis</i> f. sp. <i>tritici</i>) Septoria leaf and glume blotch (<i>Septoria</i> spp., <i>Stagonospora</i> spp.) Spot blotch (<i>Cochliobolus sativus</i>) Stem rust (<i>Puccinia graminis</i> f. sp. <i>tritici</i>) Stripe rust (<i>Puccinia striiformis</i> f. sp. <i>tritici</i>) Tan spot (Yellow leaf spot) (<i>Pyrenophora</i> spp.)	4 to 8*	2	16	Apply no later than the beginning of flowering (Feekes 10.5, Zadok's 59)
	Suppression only	6 to 8			
	Eyespot (<i>Tapesia</i> spp.)				

Application Directions. For optimal disease control, begin applications of **Priaxor** prior to disease development. To maximize yields in cereals, it is important to protect the flag leaf. Apply **Priaxor** immediately after flag leaf emergence for optimum results.

DO NOT harvest wheat hay or feed green-chopped wheat within 14 days after last application.

Priaxor does not control Fusarium head blight (head scab) or prevent the reductions in grain quality that can result from this disease. When head blight is a concern, growers should manage this disease with fungicides that are labeled for and effective in managing this disease, and with cultural practices like crop rotation and plowing to reduce crop residues that serve as an inoculum source.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 16 fl ozs of **Priaxor** per acre per season. **DO NOT** make more than two (2) sequential applications of **Priaxor** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* For early season control of Septoria leaf and glume blotch, spot blotch, and tan spot when conditions favor disease development, apply 2 to 4 fl ozs per acre of **Priaxor** either in combination with a herbicide application or when conditions favor disease development. When the 2 to 4 fl ozs early season application rate is used, a second application of **Priaxor** may be required to protect the emerged flag leaf. Environmental conditions for disease or current disease pressure at the time of flag-leaf emergence should be used to determine the **Priaxor** rate for the second application. For high disease pressure, use the higher rate of **Priaxor**.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Priaxor® Xemium® brand fungicide is specially formulated and sold by BASF for the control of pests according to the directions on this label. The purchase price of **Priaxor** includes a royalty whereby the purchaser acquires a prepaid license under U.S. Patent No. 5,438,070 and (pending No. 2008/0153707 A1) under which purchaser agrees to employ the purchased quantity of **Priaxor** only for the above-specified uses under BASF's United States patent rights and to provide notice of the terms and conditions of this license to any subsequent purchaser. Uses of **Priaxor** other than those specified on this label are not licensed through the purchase of this product and the use of this product for other purposes may violate this license and patent rights of BASF.

F500, Priaxor, and Xemium are registered trademarks of BASF.

© 2014 BASF Corporation
All rights reserved.

007969-00311.20140331b.NVA 2014-04-372-0047

Based on: NVA 2013-04-372-0017

Supplemental: NVA 2014-04-372-0095

Supplemental: NVA 2012-04-372-0100

Supplemental: NVA 2012-04-372-0125

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



We create chemistry